



The Problem with the Current High Potency THC Marijuana from the Perspective of an Addiction Psychiatrist

by Elizabeth Stuyt, MD

Advocates for the legalization of medical and retail marijuana are quick to point out all the possible benefits that a community might see from such a venture. These include increased jobs, increased tax revenue, possible medical benefits and they advertise it as “safe” and “healthy” and “organic.” They utilize the words “cannabis” and “marijuana” for everything without differentiating between the different forms of cannabis that can have very different effects on the mind and body.

Many people who have voted for legalization thought they were talking about the marijuana of the 1960s to 1980s when the THC content was less than 2%. However, without any clear guidelines or regulations from government officials, the cannabis industry has taken a page from the tobacco and alcohol industries’ play book and developed strains of marijuana and concentrated marijuana products with much higher concentrations of THC, the psychoactive component that causes addiction. The more potent a drug is, the stronger the possibility of addiction and the more likely the person will continue to purchase and use the product.



Elizabeth 'Libby' Stuyt, MD, is a board certified Addiction Psychiatrist and a Senior Instructor for the University of Colorado Health Science Program, Department of Psychiatry. She is the medical director for a 90-inpatient dual diagnosis treatment program in Pueblo, Colorado.
Contact: libbystuyt@msn.com

The active component in marijuana that people find so desirable was not really known until the 1960s when a research team in Israel found that after injecting THC into aggressive rhesus monkeys, they became calm and sedate.¹ This team discovered that there was a receptor in the brain that fit THC like a glove so they named these receptors cannabinoid receptors. It was not until the 1990s that this same team discovered why we have these receptors in our brain.¹ They discovered compounds produced by our bodies that fit into these receptors which they named anandamides, a Sanskrit word for “supreme joy.” These receptors are found all over the brain and are still called endocannabinoid receptors but that is not because they are meant for people to take in THC.

The primary problem with the current available cannabis in dispensaries in Colorado is that the THC content is not like it used to be. Prior to the 1990s it was less than 2%. In the 1990s it grew to 4%, and between 1995 and 2015 there has been a 212% increase in THC content in the marijuana flower. In 2017 the most popular strains found in dispensaries in Colorado had a range of THC content from 17-28% such as found in the popular strain named “Girl Scout Cookie.”² Sadly these plants producing high levels of THC are incapable of producing much CBD, the protective component of the plant so these strains have minimal CBD. For example the Girl Scout Cookie strain has only 0.09-0.2% CBD.

The flower or leaves that are generally smoked or vaped are only one formulation. We now have concentrated THC products such as oil, shatter, dab, and edibles that have been able to get the THC concentration upwards of 95%. There is absolutely no research that indicates this level of THC is beneficial for any medical condition. The purpose of these products is to produce a high, and the increased potency makes them potentially more dangerous and more likely to result in addiction.

Because there was initially no regulation on the edibles they have been made to look very similar to regular products that people consume such as chocolates, gummy bears, PopTarts etc. As a result there has been a significant increase in the accidental exposure/overdoses of children younger than nine in Colorado compared with the US at large.³ New regulations beginning in 2019 require that all cannabis packaging in the state of Colorado must have a universal “THC” symbol on the label with the written warning “Contains Marijuana. Keep away from Children.” All marijuana-infused products must have the universal symbol marked on at least one side of the “Standard Serving of Marijuana.”

According to the 2014 Monitoring the Future Study, marijuana is by far the number one drug abused by eighth and twelfth graders.⁴ Since legalization in Colorado, marijuana use in adolescents and those 18-25 has steadily climbed, well outpacing the national average. Colorado leads the nation in first time marijuana use by those aged 12-17, representing a 65% increase in adolescent use since legalization.⁵ According to the Colorado Department of Public Health and Environment in 2015 the county of Pueblo, Colorado, has the highest prevalence of reported past month marijuana use by high school students at 30.1%.⁶ It is well documented that when drugs are perceived as harmful, drug use decreases as we have seen with adolescent use of tobacco.⁷ There is significantly less perception of harm by marijuana primarily because Colorado has normalized it as a society and allowed the perception that it is “organic” and “healthy” and that there is nothing wrong with it.

However, there are significant consequences of long-term or heavy marijuana use beginning in adolescence. Adolescence is a time of significant brain development. Normally during this period there is a significant increase in dopaminergic and glutamatergic stimulatory neurotransmitters and a decrease in serotonergic and GABAergic suppressive neurotransmitters located in the pre-frontal motor cortex – the last part of the brain to fully develop.⁸ The prefrontal motor cortex or the “seat of judgement” is the last to fully develop and can take up to 25 – 30 years to fully develop. This equates to a great deal of learning, exploring and doing during this period, similar to stepping on the gas pedal and problems with impulse control and judgement, similar to problems stepping on the brake.

The reasons why adolescents are at such great risk for developing an addiction to drugs or alcohol is because this is a period with increased neurobiological based

tendencies for risk taking with decreased suppressive and regulatory control, and this is a period of decreased parental monitoring and increase in peer affiliations, a “perfect storm.”

The marijuana of old used to be classified as a hallucinogen and was thought to not cause addiction because there was no identified withdrawal syndrome. This has changed and with the increased potency of THC there is a definite recognized withdrawal syndrome which includes increased anger, irritability, depression, restlessness, headache, loss of appetite, insomnia and severe cravings for marijuana.⁹ It has been reported that 9% of those who experiment with marijuana will become addicted; 17% of those who start using marijuana as teenagers will become addicted; and 25-50% of those who use daily will become addicted.¹⁰ A 2015 study carried out in the UK found that high-potency cannabis use is associated with increased severity of dependence, especially in young people.¹¹

Addiction is a problem with the learning and memory part of the brain and all drugs of abuse work in the same “reward pathway” where we learn to do anything such as eat and procreate. All drugs of abuse cause a release of dopamine from the nucleus acumbens that signifies salience and starts the process of long term potentiation which reinforces the learning. At the same time, the hippocampus which is vitally important for new memory and learning is negatively impacted by the chronic use of any addictive substance. These substances decrease neurogenesis in the hippocampus and actually cause shrinkage of the hippocampus and impair the ability to learn new things. This is true for alcohol, cocaine, methamphetamine, heroin, nicotine, and THC.¹² Animal studies have demonstrated impaired learning with all of these substances but the good news is that recovery is possible. When the use of addictive drugs is stopped and the animals are allowed to be in a recovery environment where they are free to exercise (voluntary exercise being one thing that improves neurogenesis) they can again learn new things.¹³

Human studies have shown that long-term (> 10 years) and heavy (> 5 joints per day) cannabis use compared with age matched non-using controls resulted in bilaterally reduced hippocampal and amygdala volumes ($p=.001$) and significantly worse performance on measures of verbal learning ($p<.001$).¹⁴ There is evidence that recovery is possible in humans as well. A study of 40 male and 34 female long-term (@15 years) cannabis users versus 37 non-users, healthy controls divided the marijuana users into three groups; those that smoked predominantly THC in the previous three months, those who smoked a combination

PERSPECTIVE

of THC and CBD in the previous three months and former users with a sustained abstinence of 29 months.¹⁵ They found that cannabis users had smaller hippocampal volumes compared to controls but the users not exposed to CBD had an even greater (11%) reduced volumes (CBD appears to be somewhat protective). In the former users the hippocampal integrity was comparable to controls. The only problem with this study is they did not test for functional deficits to see if function improved along with hippocampal volume.

There are other important neurotransmitters that are very active during adolescence and include acetylcholine receptors (ACH) and endocannabinergic receptors (CB1). ACH helps us focus and concentrate and ACH innervation of the pre-frontal motor cortex reaches mature levels during adolescence.¹⁶ These receptors in the brain are called nicotinic or nAChRs to differentiate them from the muscarinic receptors in the body. They are called nicotinic simply because nicotine binds to these receptors – not because we are supposed to use tobacco products. These receptors are involved in promoting or preventing neuronal cell death depending on the stage of brain development. Putting an exogenous form of nicotine in the developing brain, as in consuming tobacco, can dysregulate these fine tuning mechanisms during adolescence.

CB1 receptors regulate the balance between excitatory and inhibitory neuronal activity utilizing our own natural anandamides. Exposure to cannabis during adolescence disrupts glutamate which plays an important role in synaptic pruning in the pre-frontal motor cortex; disrupting normal brain development.¹⁷ This is most likely why there are many studies demonstrating the negative effect on cognition and IQ in people who are exposed to marijuana beginning in utero through adolescence. In spite of this, nearly 70% of dispensaries in Denver, Colorado, recommend cannabis products to treat nausea in the first trimester of pregnancy.¹⁸ This is basically bud-tenders practicing medicine without a license.

A study in New Zealand with a 20-year follow-up showed an average loss of 8 IQ points with early persistent teen use of marijuana.¹⁹ If you already have a high IQ, a drop in 8 points may mean the difference between making As and making Bs, however for the person with an average IQ of 100 (50th percentile), a loss of 8 points can put that person in the 29th percentile with significant difficulty in functioning. A study out of Yale University tracked 1,142 students who achieved similar SAT scores and were enrolled in college.²⁰ They found that those who used minimal alcohol or cannabis had an average GPA of 3.1 at the end

of the semester. Those who drank alcohol without using marijuana had an average GPA of 3.03 and those who used both alcohol and marijuana had an average GPA of 2.66.

Marijuana use is also correlated with creating or worsening many mental health problems including anxiety, depression, psychosis, and suicidal ideation. A prospective study in Australia followed 1,600 girls for seven years starting before they expressed symptoms of mental illness or substance abuse.²¹ They found that girls who used marijuana at least once a week were twice as likely to develop depression than those who did not use, and those who used marijuana every day were five times more likely to suffer from depression and anxiety than non-users. A study of 307 adults with depression assessed symptoms, functioning and marijuana use at baseline, and three- and six-month intervals.²² Researchers found that marijuana use was associated with poor recovery. Those aged 50+ increased their marijuana use compared to the youngest age group ($p < .001$) and the marijuana use worsened depression ($p < .001$) and anxiety ($p = .025$) symptoms. Marijuana use led to poorer mental health functioning compared to those who did not use marijuana ($p = .01$).


Numerous studies have demonstrated that using cannabis prior to the age of 15-18 significantly increases the risk of developing psychotic symptoms.²³ The risk is dose dependent and increases with greater frequency of use and with higher potency THC. A landmark study out of the UK analyzed 780 adults, ages 18-65, 410 with their first psychotic episode versus 370 matched healthy controls.²⁴ They found that use of high potency THC > 15% resulted in a three times increased risk of psychosis, and if the use was daily there was a five times increased risk. Those using hash with < 5% THC did not exhibit psychotic symptoms.

A growing number of states have identified PTSD as an approved condition for medical marijuana. However, this is not based on any research. There is no evidence that marijuana successfully treats PTSD and there is evidence that it can make it worse. Marijuana is not the answer for PTSD similar to the reason why benzodiazepines or alcohol are not the answer for PTSD. All these compounds do is provide temporary relief by numbing the individual and disconnecting them from the traumatic emotion. It does not resolve the trauma, and they have to continue to use multiple times a day in order to continue with the benefit. This can lead to increased addiction potential and withdrawal symptoms, cognitive impairment, a-motivational syndrome, and the potential for psychosis or worsening psychosis from the PTSD. An observational study done by the VA followed 2,276 Veterans who were treated for PTSD

in one of the VA PTSD treatment programs around the country.²⁵ It compared those using marijuana and those not using it and found those who never used marijuana had significantly lower symptom severity four months after PTSD treatment. Those who were using marijuana but stopped using it in treatment had the lowest level of PTSD symptoms four months after treatment, and those who started smoking marijuana had the highest levels of violent behavior and PTSD symptoms four months after treatment. Another conundrum that impacts treatment for PTSD is the possibility that cannabis users have an increased susceptibility to memory distortions even when abstinent and drug free which can compromise reality monitoring. Riba et al. studied 16 heavy cannabis users (daily for last two years – average of 21 years) to 16 matched cannabis naïve controls.²⁶ The cannabis users had to abstain from cannabis use for four weeks prior to the study. The study involved a memory paradigm including a study phase and a testing phase with the participant in an MRI scanner. They were given lists of four words to memorize and then shown a different list and they had to report if the words were on the previous list. Marijuana users were significantly more likely to have false recognition of the words and were less likely to reject that they had a false memory compared with the non-users.

Multiple studies have documented a relationship between cannabis use and suicidality. A large, longitudinal study in Australia and New Zealand of over 2000 adolescents and maximum frequency of marijuana use found almost a seven fold increase in suicide attempts in daily marijuana users compared with non-users.²⁷ A Congressional Hearing on April 27, 2017, reported that Veteran suicides were up 32% since 2001 compared to a national increase of 23% during the same time period. A 2017 cross-sectional multi-site VA study of 3,233 Veterans found that cannabis use disorder was significantly associated with both current suicidal ideation ($p < .0001$) and lifetime history of suicide attempts ($p < .0001$) compared to Veterans with no lifetime history of cannabis use disorder.²⁸ This significant difference continued even after adjusting for sex, PTSD, depression, alcohol use disorder, non-cannabis drug use disorder, history of childhood sexual abuse, and combat exposure. According to the Colorado Department of Public Health and Environment, marijuana is by far the most frequently encountered drug on toxicology screens of suicides among adolescents ages 10 – 19 and has been increasing over the last eight years.²⁹

Misguided marijuana advocates have recently been suggesting that marijuana is a solution for the opioid epidemic. There is no clinical evidence of this and in fact, marijuana is found to be more of a “companion” drug rather than an “alternative” drug for most patients seeking addiction treatment in Colorado. A study of 5,315 adolescents in the UK with three or more measures of cannabis use from age 13-18 found a dose-response relationship between cannabis use trajectories in adolescence and nicotine dependence, harmful alcohol consumption, and other illicit drug use by age 21.³⁰ A large study of 34,653 individuals using NESARC data compared cannabis use at wave 1 (2001-2002 – 81% response rate) to prescription opioid use disorder at wave 2 (2004-2005 – 70.2% response rate).³¹ Cannabis use at wave 1 was associated with a significant increase of having a prescription opioid use disorder at wave 2, with over four times the risk for those who had frequent use of marijuana.



There is evidence that prenatal exposure of cannabis can alter opioid gene function in humans. Fetal brains obtained from aborted fetuses from women who were using marijuana during their pregnancy were compared to those from women not using marijuana during pregnancy.³² The researchers discovered impaired opioid-related genes in distinct brain circuits that they hypothesized may have long term effects on cognitive and emotional behavior. These findings are comparable to findings with animals. One study of prenatal cannabis exposure in rats found that the THC exposed rats exhibited shorter latency to first active lever press for heroin and had higher heroin-seeking during mild stress and drug extinction than animals not exposed to THC.³³ The THC exposed animals exhibited allostatic changes in the limbic encephalin systems in adulthood.

Another interesting study that supports the idea that cannabis use and opioid use are linked was in a randomized, double-blind, placebo controlled trial of naltrexone in non-treatment seeking cannabis smokers.³⁴ In a laboratory setting those receiving a placebo had 7.6 times the odds of self-administering active cannabis compared with those receiving daily naltrexone, an opioid receptor blocker.

If states continue to commercialize marijuana as has been done in Colorado we are destined to see many more people requiring treatment for addiction, depression, anxiety, suicidal ideation, and psychosis. We need to continually educate every one of the risks and increase prevention efforts to prevent children and adolescents from initiating marijuana use. This should include a strong ban on any advertising that appears to be directed toward youth – for all drugs including marijuana, tobacco, and alcohol. States will need to commit to increased funding for and availability of treatment options. The strongest recommendation would be to initiate regulations to limit the concentration of THC. Ideally this would be to less than 10% as there is no good research on concentrations greater than this for any medical condition and there is significant literature on the negative effects of high potency THC.

References

- Sides H. Science seeks to unlock marijuana secrets. National Geographic Magazine. June 2015
- www.leafly.com – accessed July 15, 2017
- Wang GS et al. Unintentional pediatric exposures to marijuana in Colorado, 2009–2015. *JAMA Pediatr.* 2016;170(9):e160971.
- University of Michigan, 2014 Monitoring the Future Study <http://www.monitoringthefuture.org/pubs/monographs/mtf-overview2014.pdf>
- Rocky Mountain High Intensity Drug Trafficking Area. Marijuana in Colorado: the impact. 2017;5 <http://www.rmhidta.org/html/FINAL%202017%20Legalization%20of%20Marijuana%20in%20Colorado%20The%20Impact.pdf>
- Colorado Department of Public Health and Environment. Healthy Kids Survey. 2015 https://www.colorado.gov/pacific/sites/default/files/PF_Youth_HKCS_MJ-Infographic-Digital.pdf
- University of Michigan, 2013 Monitoring the Future Survey. NIDA <https://www.drugabuse.gov/publications/drugfacts/monitoring-future-survey-high-school-youth-trends>
- Schepis et al. Neurobiological Processes in Adolescent Addictive Disorders. *Am J Addictions.* 2008;17:6-23
- Bonnet U, Preuss UW. The cannabis withdrawal syndrome: current insights. *Sub Abuse Rehab* 2017;8:9-37.
- Volkow ND et al. Adverse Health Effects of Marijuana Use. *N Engl J Med* 2014;370:2219-2227.
- Freeman TP; Winstock AR. Examining the profile of high-potency cannabis and its association with severity of cannabis dependence. *Psychol Med* 2015;45:3181-3189.
- Chambers RA. Adult hippocampal neurogenesis in the pathogenesis of addiction and dual diagnosis disorders. *Drug Alcohol Depend* 2013;130:1-12.
- Mandyam CD, Koob GF. The addicted brain craves new neurons: putative role for adult-born progenitors in promoting recovery. *Trends Neurosci* 2012;35:250-260.
- Regional brain abnormalities associated with long-term heavy cannabis use. *Arch Gen Psychiatry* 2008;65:694-701.
- Yucel et al. Hippocampal harms, protection and recovery following regular cannabis use. *Transl Psychiatry* 2016;6:e710
- deBry SC, Tiffany ST. Tobacco-induced neurotoxicity of adolescent cognitive development (TINACD): A proposed model for the development of impulsivity in nicotine dependence. *Nicotine & Tobacco Research* 2008; 10:11-25.
- Lubman et al. Cannabis and adolescent brain development. *Pharmacology and Therapeutics* 2015;148:1-16
- Dickson B et al. Recommendations from cannabis dispensaries about first-trimester cannabis use. *Obstetrics and Gynecology* 2018;131:1031-1038.
- Meier MH et al. Persistent cannabis users show neuropsychological decline from childhood to midlife. *PNAS* 2012; E2657-E2664. doi/10.1073/pnas.1206820109
- Meda SA et al. Longitudinal influence of alcohol and marijuana use on academic performance in college students *PLOS ONE* | DOI:10.1371/journal.pone.0172213 March 8, 2017
- Patton GC et al. Cannabis use and mental health in young people: cohort study. *BMJ*;2002;325:1195-1198.
- Bahorik AL et al. Patterns of marijuana use among psychiatry patients with depression and its impact on recovery. *J Affect Disord.* 2017; 2013:168-171.
- Pierre JM Risks of increasingly potent Cannabis: the joint effects of potency and frequency. *Current Psychiatry* 2017;16:14-20
- DiForti et al. Proportion of patients in south London with first-episode psychosis attributable to use of high potency cannabis : a case-control study. *Lancet Psychiatry*, 2015; doi.org/10.1016/S2215-0366(14)00117-5.
- Wilkinson et al. Marijuana use is associated with worse outcomes in symptom severity and violent behavior in patients with posttraumatic stress disorder. *J Clin Psychology* 2015;76:9.
- Riba et al. Telling true from false: cannabis users show increased susceptibility to false memories. *Molecular Psychiatry* 2015;20:772-777.
- Silins E et al. Young adult sequelae of adolescent cannabis use: an integrative analysis. *Lancet Psychiatry* 2014;1:286-293.
- Kimbrel NA et al. Cannabis use disorder and suicide attempts in Iraq/Afghanistan-era veterans *J Psychiatric Research* 2017;89:1-5.
- https://www.colorado.gov/pacific/sites/default/files/CHED_VS_Health-Watch-No-94-Adolescent-Suicide-in-Colorado-2008-2012_0817.pdf
- Taylor M et al. Patterns of cannabis use during adolescence and their association with harmful substance use behavior: findings from a UK birth cohort. *J Epidemiol Community Health.* 2017; 0:1–7. doi:10.1136/jech-2016-208503.
- Olfson M et al. Cannabis use and risk of prescription opioid use disorder in the United States. *AJP in Advance* (doi:10.1176/appi.ajp.2017.17040413).
- Wang et al. Prenatal exposure of cannabis alters opioid gene function in humans. *Pharmacogenomics J*, 2006;6:255-264.
- Sapano et al. Prenatal cannabis exposure increases heroin seeking in adult rats. *Biol Psychiatry* 2007;61:554-563.
- Haney et al. Naltrexone maintenance decreases cannabis self-administration and subjective effects of daily cannabis use. *Neuropsychopharmacology* 2015.



FUNDING: Dr. MacDonald's work is sponsored in part by the Goodenough Neuroscience Research Fund.

FINANCIAL DISCLOSURES: The authors have no conflicts of interest relevant to the content of this article.

ADDRESS CORRESPONDENCE TO:

Kai MacDonald, MD, 3368 2nd Avenue, Suite B, San Diego, CA 92103; Phone: (619) 203-7393; Fax: (619) 296-0199; E-mail: kai@kaimacdonald.com

KEY WORDS: Marijuana, cannabis, dependence, drug-related harms, mental health

REVIEW

WHY NOT POT?

A Review of the Brain-based Risks of Cannabis

by **KAI MACDONALD, MD, and KATHERINE PAPPAS, BA**

Dr. MacDonald and Ms. Pappas are with UC San Diego Psychiatry, San Diego, CA, USA

Innov Clin Neurosci. 2016;13(3-4):13-22

ABSTRACT

In this review, we provide a historical perspective on marijuana, and survey contemporary research investigating its potential negative effects on the brain. We discuss the evidence regarding cannabis dependence, driving under the influence of cannabis, underachievement, inducing (or worsening) certain psychiatric conditions, and the potential for progression to use of more dangerous drugs—summarized by the acronym DDUMB, a cognitive tool that may help healthcare providers in their risk/benefit discussions with patients who use cannabis. We also review and discuss the impact of marijuana use on target populations, including adolescents (who are at increased risk of harm); heavy users; and people suffering from—or at high risk of—mental illness. While cannabis presents certain subjective, health-related, and pecuniary benefits to users, growers, and other entities, it is also associated with several brain-based risks. Understanding these risks aids clinicians and their patients in making informed and balanced decisions regarding the initiation or continuance of marijuana use.

INTRODUCTION

“I have argued that every human being is born with an innate drive to experience altered states of consciousness periodically . . . this drive is a most important factor in our evolution, both as individuals and as a species.”

Andrew Weil

The Natural Mind: A Revolutionary Approach to the Drug Problem

Marijuana, also known as cannabis or pot, is the most commonly used drug worldwide and is a fraught topic in contemporary society.¹ A variety of forces—economic,² legislative,³ technological,⁴ and even horticultural⁵—have markedly changed the politics, polemics, and public perception of pot. The resultant upsurge of cannabis use in some parts of the United States has already had a collateral impact on individual and societal health,⁶ similar to that seen with the prescription opiate epidemic.⁷ Balancing these myriad forces—all of which drive greater societal acceptance of marijuana and increased use—a growing body of

scientific research provides a clearer understanding of pot's potential harms.

The aim of this paper is to review the brain-based harms of cannabis. Awareness of the supporting evidence of marijuana's downsides can help augment the risk-benefit discussions clinicians may have with patients in a motivational interviewing model, the preferred therapeutic frame for approaching substance-use discussions.⁸ To facilitate this end, we introduce a mnemonic, DDUMB, to help remind us of the five brain-based harms associated with marijuana use: dependence, driving impairment, underachievement, mental illness, and bad to worse (i.e., marijuana serving as a "gateway" function for other more dangerous drugs of abuse). Before reviewing the science behind these five dangers, we will provide a brief summary of several important aspects of marijuana's history, politics, chemistry, and psychopharmacology.

THE HISTORY OF HEMP AND POLITICS OF POT

The terms *marijuana* and *cannabis* are often used interchangeably. Strictly speaking, however, *cannabis* is a botanical term for the hemp plant, while *marijuana* denotes the psychoactive drug derived from it. Though research on the central effects of cannabis is relatively new, its medicinal use can be traced back to the Chinese Han dynasty, circa AD 25 to AD 220, when it was used to treat rheumatic pain, constipation, malaria, and female reproductive disorders.⁹ Medical cannabis was introduced to the Western world in the 1800s, and was used as such until the 1900s, when its popularity diminished with the rise of pharmaceuticals that could be used for the same conditions (e.g., aspirin, barbiturates).¹⁰ Legislation enacted in 1937 (the Marijuana Tax Act) decreased accessibility and pushed the drug further out of the public eye. A sterling demonstration of the swings of public opinion toward

marijuana use is the pulp-propaganda film "Reefer Madness,"¹¹ released shortly after the enactment of this legislation. Originally titled "Tell Your Children," this short film comically overdramatized marijuana's harms, describing cannabis as "the burning weed with its roots in Hell," and warning about the potential for pot-induced manslaughter, suicide, hallucinations, and "the ultimate end of the marijuana addict: hopeless insanity!"

In the present day, more tolerant state-based legislation has led to decriminalization, legalization, and medicalization of cannabis in many states and the likelihood for more cannabis use. In turn, more frequent use by current users and more new users may lead to a greater frequency of cannabis-related harms. Specifically, in December of 2012, Washington state and Colorado both legalized marijuana; Washington DC, Alaska, and Oregon followed suit, with at least 23 states now allowing for its medical use. Importantly, this large cultural and legal pro-pot shift has already been shown to increase a variety of cannabis-related collateral harms,⁶ and likely has contributed to an increase of adolescent-onset use.¹² This trend is especially worrisome, since adolescent-onset use is associated with greater cumulative negative consequences than later-onset use.¹³ Bachman et al¹³ demonstrated an inverse relationship between perceived risks/social disapproval and the prevalence of marijuana use among youth.¹³ Data from a 2010 national survey on drug use and health have shown a correlation between adolescent cannabis use and lower levels of parental disapproval.¹⁴ Additionally, "vaping," a term used to describe a popular method of smoking via an electronic device such as an e-cigarette, may encourage more illicit and dangerous use of marijuana: Vaping produces less smoke than marijuana or tobacco cigarettes, making its use harder to detect by smell (e.g., in a school bathroom) and implies that the person is vaping

nicotine-related products (not illicit substances).¹⁵ These relationships suggest a continuing trend toward public approval and, possibly, risk-minimization of marijuana use, which may lead more at-risk youth and young adults to initiate use.

Unfettered by more permissive laws and attitudes, capitalism has embraced cannabis as the newest cash crop.^{16,17} Commercialization of cannabis has been shown to increase the number of medical marijuana licenses purchased.²¹ Therewith, powerful economic forces have and likely will continue to add to legal and attitudinal shifts that elevate the role of cannabis in public and individual health.²³ Market research suggests that widespread legalization of marijuana has the potential to create a 35-billion dollar marijuana market.¹⁸ Comparatively, this would make the marijuana industry as big as the United States National Football League (NFL), 10 times more profitable than the opioid drug OxyContin, and about a fifth the size of the United States alcohol market.^{18,19} As witnessed in recent years in the United States, where a surge in opiate-related mortality has been partly attributed to high-dollar opiate sales, the promise of profits in the burgeoning industry of marijuana production—the "green" industry—may inform how the drug is marketed and researched.^{7,20} And as more states move to legalize marijuana, we may see an increase in both anticipated and unanticipated cannabis-related harms in those who use it, as was observed on a smaller scale in Colorado.^{6,21}

As societal, legislative, and economic forces move toward the legalization of marijuana, there are three challenges that confront its scientific study. First, advances in cultivation techniques and grower knowledge have produced vastly more potent marijuana than was seen in previous decades. Tetrahydrocannabinol (THC), one of the main psychoactive components of marijuana (and the component associated with some of its brain-

based harms⁷⁴) has increased in concentration from three percent in the 1980s to 12 percent in 2014, whereas the concentrations of cannabidiol (CBD), one of marijuana's calming components, has fallen.⁷ This horticultural reality makes older literature on the effects of marijuana less applicable to current use, and is frequently cited as a reason for a relative increase in cannabis-related harms.^{6,23}

Aside from differences in the chemical composition of the plant, a second, related challenge in marijuana research is that precise quantification of cannabis use (compared to a drug like nicotine found in tobacco) is difficult, due to differing potencies and variable delivery systems (e.g., smoking, ingestion). A third and final factor that creates challenges for cannabis research is its United States Drug Enforcement Agency (DEA) scheduling. Because marijuana is grouped with cocaine, heroin, and 3,4-Methylenedioxymethamphetamine (MDMA) in the most restrictive drug schedule (Schedule 1), access to the drug for scientific study is more difficult. From the perspective of advancing scientific knowledge, many researchers have suggested moving marijuana to a lower schedule to reduce barriers to research.^{24,25}

CANNABIS CHEMISTRY AND PHARMACOLOGY

Unlike drugs that contain a single, specific, active chemical compound (e.g., lithium), different strains of the cannabis plant contain produce a variable array of centrally active substances. One of the main chemical groups in the several dozen constituents of marijuana are the cannabinoids, which become active by binding at cannabinoid receptors in the human brain. The three primary cannabinoids—found in varying ratios in different strains of cannabis—are cannabitol (CBN), cannabidiol (CBD), and Delta-9-tetrahydrocannabinol (THC). Importantly, CBD and THC often have opposing effects. CBD has anxiolytic and antipsychotic properties, and is

often marketed as such, while THC has been shown to be anxiogenic and can induce transient psychosis.⁵ From this chemical complexity follows the clinical reality discussed above: different strains of the cannabis plant, since they contain different ratios of centrally active chemicals, yield different central effects. This “blending”—rooted in the inherently variable chemistry of plant-based drugs—yields a compound that has different effects and potential harms in each of its permutations. These factors add additional challenge to the scientific study of marijuana, given that the actual drug one is studying may vary depending on the several variables (e.g., when it was grown, where it was grown).

Though marijuana has been around for millennia, our understanding of its mechanism of action in the brain is relatively recent. This understanding was propelled forward in the late 1980s by the discovery of central cannabinoid receptors, which bind both marijuana-derived cannabinoids as well as the brain-made substances called endogenous cannabinoids (or endocannabinoids).⁵ Two sets of cannabinoid receptors, called CB1R and CB2R, exist. CB1R is located in widespread brain regions (i.e., hippocampus, cerebral cortex, limbic system, cerebellum), but is also found in the periphery (i.e., in liver, thyroid, uterus, bones, and testicular tissue). CB2R, on the other hand, is mainly peripheral, found mostly in immune cells, the spleen, and gastrointestinal system.⁵

Comparing endogenous and exogenous cannabinoids is informative. The primary endogenous endocannabinoid, anandamide, was named from the Sanskrit word for “supreme joy.”¹⁰ Unlike THC, which has a half-life that spans hours to days,²⁶ anandamide has a short half-life, being quickly removed from the synapse and degraded.¹⁰ The difference in half-lives between anandamide and THC may contribute to some of the different central effects of these two molecules,

including the potential for dependence.²⁷

DEPENDENCE

The first cannabis-related harm captured in the DDUMB mnemonic is dependence. Substance dependence is a debilitating, brain-based disorder characterized by compulsive use, inability to desist in the face of negative consequences, and withdrawal symptoms upon cessation.²⁸ Although commonly believed to be nonaddictive, marijuana dependence has been clearly documented,²⁹ and a large percentage of global substance use admissions are related to cannabis.³⁰ Very recent prevalence data in the United States indicate that past-year prevalence of marijuana use doubled between the years 2001 and 2013 to nearly 10 percent, with a corresponding increase in marijuana use disorders to nearly three percent.³¹ Though neurobiological responses are not yet used to validate substance dependence, THC has been shown to stimulate mesolimbic dopamine release, a brain phenomenon common to all addictive substances,²⁸ and many other experiments indicate that cannabis affects key parts of the brain's addiction centers.³²⁻³⁴

The misperception that cannabis is not addictive has at least three sources. First, the percentage of first-time cannabis users who develop dependence is relatively low compared to other commonly abused drugs. Specifically, nine percent of first-time cannabis users get hooked versus higher percentages of first-time stimulant (11%), alcohol (15%), cocaine (17%), heroin (23%), and nicotine (32%) users.³⁵ That said, though the percentage of first-time cannabis users who develop dependence is lowest among users of the other drugs mentioned, the overall number of people who will develop cannabis addiction is still large.

A second factor supporting the non-addiction myth is that although chronic cannabis users typically

“dose” multiple times a day, the long half-life of THC (25–57 hours)²⁶ means that the time intervals that mark its “compulsive use” can be spaced out longer than other shorter-acting drugs, such as nicotine, creating more of an illusion of control. The addicted marijuana user may only use pot at breakfast, lunch, and in the evening, whereas a person addicted to nicotine may need to smoke a cigarette every hour or two.

A third factor supporting the myth of non-addiction is that marijuana withdrawal is often relatively mild.^{27,29} Moreover, marijuana withdrawal presents without clear, “signature” physical symptoms, at least compared to the often-dramatic physical symptoms of withdrawal from depressants like alcohol and benzodiazepines (i.e., tremors, seizures, agitation) and opiates (i.e., sweating, gooseflesh, diarrhea). Instead, marijuana withdrawal symptoms are more occult: anorexia, irritability, anxiety, anger, restlessness, and sleep disruption.¹³ En toto, this delayed, nondramatic withdrawal syndrome adds support to the misperception that cannabis addiction does not exist.

For clinicians treating marijuana-addicted patients, it is important to be aware that cannabis withdrawal is both consequential and treatable.^{2,37} In terms of its consequence, cannabis withdrawal symptoms clearly contribute to ongoing use, making cessation efforts aversive,^{28,38,39} and to impairing both motivation and executive functions critical in decision making and treatment retention.^{40,41} In terms of treatment, a seminal study by Mason et al³⁷ showed that in marijuana-dependent subjects, the commonly used calcium channel/GABA-modulating agent gabapentin—dosed 1200mg daily in divided doses—reduced both cannabis withdrawal symptoms and cannabis use. Though sustained recovery from cannabis addiction requires long-term, multimodal solutions, clinicians can help the process by utilizing available pharmacotherapies to attenuate withdrawal.

It is important to highlight that the risk of negative effects from marijuana use—including dependence—have been shown to be related to the age of first exposure.^{42,43} Specifically, compared to people who start marijuana use in adulthood, adolescent initiators are 2 to 4 times more likely to exhibit dependence within two years of their first use.⁴² This is not surprising, given that key stress, reward, and executive/regulatory circuits that underlie addiction continue to develop during the teenage and early adult years of human growth.^{28,44} Moreover, studies indicate that early exposure to THC may 1) potentiate the future effects of THC, increasing risk of dependence; 2) cause impaired regional connectivity, decreasing the moderating influence of regulatory brain regions; and 3) lead to lower dopaminergic activity in addiction-related circuits.^{23,35,46} Coming from the perspective of harm reduction, then, a tractable goal to reduce the risk of future marijuana dependence is to delay the age of onset of first use.

DRIVING

Standing alongside the misconception that marijuana is not addictive is the misconception that driving while under the influence of marijuana is safe. Several factors make this latter untruth more challenging to refute than the former. Until very recently, drivers involved in accidents or infractions were rarely tested for THC levels, whereas assessing blood alcohol content via the less invasive breathalyzer has been routine for years. This situation will likely change over time as marijuana use increases, allowing more THC-related auto morbidity studies to be conducted and compared with those in other countries. A second factor complicating the THC-driving research is that, based on studies from driving fatalities, drivers frequently use marijuana and alcohol simultaneously.⁴⁵ This combination makes assigning causality to a single

drug complex, and their different mechanisms of action lead to additive impairments.⁴⁷

Despite the abovementioned challenges to studying the topic, epidemiological and laboratory studies of the acute effects of marijuana on driving has demonstrated that drivers under the influence of marijuana are impaired. In fact, driving while under the influence of marijuana doubles or triples the risk of a crash.⁴⁷ Though people driving under the influence of marijuana tend to compensate by driving more slowly, as task intensity of driving increases, the person becomes more impaired.^{48,49} Specifically, cannabis use increases lane weaving and impairs critical-tracking tasks, reaction time, and divided attention.^{48,49}

Though a discussion of the ethical issues of driving while impaired is beyond the scope of this article, it bears mentioning that collateral damages result from individual choice—every time an impaired motorist decides to get behind the wheel, he or she extends the risk of potential harms to other drivers, passengers, pedestrians, and cyclists.

UNDERACHIEVEMENT

Despite marijuana’s known risks, the scientific reality is that marijuana is in many ways the least deadly drug of abuse. In meteorological terms, if methamphetamine—with its capacity for brain damage and strokes⁵⁰—is a tsunami, and opioids—with their morbid respiratory depressive effects⁷—are an earthquake, marijuana can be likened to a heavy fog. Disruptive, yes. Deadly, no. Partly due to the lack of activity in vital brainstem areas controlling respiration, there has never been a reported lethal overdose of marijuana in humans.⁵¹ In animals, the deadly dose of cannabis is extremely high: about 12,500 times the amount needed to cause subjective effects.⁵² Though risks of marijuana use are real and consequential, it is neither deadly nor overly dramatic. In the pot polemic, the lack of direct organ

toxicity, clearly consequent mortality, and extreme withdrawal symptoms likely contribute to the growing acceptance of marijuana use among the American population. And as argued in a recent editorial by a pediatrician in the *New York Times*, support for marijuana use may come from the perspective, “Since people are going to use something, why not the least toxic something?”⁵³

From a wider social perspective of harm and risk reduction, this “low bar” argument has obvious merits: it is better to be alive and stoned than dead from a heroin or alprazolam overdose. That said, conscious of Weil’s opening observation about our species’ “innate drive to experience altered states of consciousness,” we suggest that when considering effects on vulnerable future adults, professionals should focus not on what is the least toxic and not on merely accepting morbidity over mortality, but rather on the maximization and optimal development of human potential. From this vantage, the risk of cannabis promoting broad-spectrum underachievement (especially in teens) becomes more prominent. In point of fact, a large body of convergent data suggest that long-term use of marijuana may cause significant abridgement of one’s potential.^{33,44,54–59,60–67}

Underachievement may be the most well-supported correlate of regular marijuana use. Though direct causality is challenging to ascribe due to the correlative nature of this research (random assignment of daily cannabis use to adolescents is unethical), the association of daily marijuana use with the pruning of human potential appears across a breadth of contemporary research. For example, earlier studies have already demonstrated that marijuana use during adolescence is associated with low academic achievement and increased rates of school drop-out.^{54,55} More recently, several very large examinations of the issue have reinforced not only these academic consequences, but a broader swath of

negative outcomes. In one of these studies, Fergusson et al⁵⁶ performed a longitudinal study of over 1,000 New Zealanders from birth to age 25 years, and found that elevated marijuana use between ages 14 and 21 years was associated with the lower likelihood of getting a bachelor’s degree, lower income, higher unemployment and welfare dependence, and lower levels of relationship and life satisfaction. These correlations survived adjustments for a number of important covariates, including family socioeconomic status, maltreatment, academic achievement, and comorbid mental disorders. In a similar study, Meier et al⁵⁷ followed a cohort of 1,037 subjects from birth until age 38 years, performing neuropsychological assessments at ages 13 and 38 years, as well as ascertaining cannabis use at ages 18, 21, 26, 32, and 38 years. In this cohort, persistent cannabis use was associated with a decline in neuropsychological performance across domains, which survived controlling for years of education. Importantly, these results were the most prominent among participants with adolescent-onset cannabis use, and showed a dose effect: more persistent use was associated with a more severe performance decline. Adolescent-onset cannabis use was correlated with a 10-point decrease in measured IQ. Moreover, people who had discontinued cannabis use did not achieve a full return to their baseline level of performance, a finding which suggests that heavy adolescent-onset cannabis use may have a cumulative neurotoxic effect. One group of detractors argued that certain brain-based personality traits that bias people toward marijuana use as well as school dropout may explain these results,⁵⁸ but the original authors’ results survived a control for such personality factors.⁵⁹

What are the putative mechanisms wherein adolescent cannabis use causes this pervasive underachievement and even cognitive decline? Adolescence, we know, is a neurodevelopmental stage of significant import in which

neurobiological circuits critical to adult function develop, are pruned, and reinforced.⁶⁰ Moreover, adolescent brains have a stage-dependent hypersensitivity to rewards²³ and underdeveloped prefrontal inhibitory structures.⁴⁴ Chronic cannabis use through this sensitive window of development may cause persistent disruptions in these developing prefrontal and reward pathways, impacting important intellectual functions like working memory, sustained attention, verbal memory, and general intellectual functioning.^{61–63} These disruptions may persist longer—and the person may not fully recover—when experienced in the developmental window of adolescence rather than in adulthood. Aside from prefrontal cortex disruptions, chronic marijuana use has also been correlated with changes in the hippocampus, a vital brain structure involved with short-term memory, long-term memory, and spatial navigation.⁶⁴

Recent research on the effects of marijuana on brain function and structure (in both adolescents and adults) have shown other negative changes in the brain among chronic marijuana users.^{65–67,35} In a study that compared chronic marijuana users with non-using adults (mean age 22–23 years), the chronic user group demonstrated poorer learning from errors, due in particular to lower levels of brain activity in the dorsal anterior cingulate cortex and hippocampus.⁶⁵ In another study, investigators found microscopic disturbances in the neural fibers that communicate between brain hemispheres (the corpus callosum) in heavy cannabis users (mean age 30 years) who started using at the age of 16 years.⁶⁶ In a 25-year follow-up study, investigators compared three domains of cognitive function (verbal memory, processing speed, and executive function) in three groups (aged 18–30 years at baseline): 1) current cannabis users, 2) individuals who used marijuana but stopped, and 3) individuals who never used marijuana. The researchers found that

current marijuana users demonstrated lower verbal memory and processing speed compared to the other groups, and even when current users were excluded, cumulative exposure was associated with worse verbal memory.⁶⁷ And in a study that examined brain functioning among chronic marijuana users (aged 21–33 years), investigators found impairments in dopamine release in the striatum that correlates with deficits in neurocognitive performance (memory, attention).³⁵

In summary, broad-spectrum, lasting underachievement—perhaps mediated by disruptions of critical developmental brain circuits—is a third potential harm from cannabis. Convergent evidence from several fields, including epidemiology and functional brain imaging, supports the idea that one of the more occult (but consequential) downsides of adolescent-onset marijuana use is a broad-spectrum abridgement of human potential.

MENTAL ILLNESS

Marijuana use has been associated with several specific brain-based illnesses. Much of this research has focused on the role of cannabis in psychotic illness. Though the details of this research arena are beyond the scope of this article (see references 64, 68, and 69 for more comprehensive treatment of this issue), the emerging theory follows a stress-diathesis model and posits that in genetically “at risk” individuals, marijuana use serves as a biological trigger that influences the full expression of what otherwise may have been a latent disorder. This body of research suffers the same shortcomings noted in the underachievement section: random assignment to the experimental condition—heavy cannabis use—is unethical. That said, a raft of studies have found strong support for a pot-psychosis link, indicating that cannabis use can increase the risk for the development of psychotic disorders⁶⁹ and worsen clinical outcomes in those at risk.⁷⁰ In a 35-

year longitudinal study of more than 50,000 enlisted men, Manrique-Garcia et al⁶⁸ found that individuals who used cannabis frequently had an increased long-term risk for developing schizophrenia, whereas the risk declined for moderate users.

Assessing the genetics of at-risk individuals, Caspi et al⁷¹ reported that adolescent cannabis users carrying a permutation of the catechol-O-methyltransferase (COMT) gene were at highest risk of developing psychotic illness. Mechanistically, THC increases task-irrelevant neural “noise,” which is associated with its psychosis-promoting effects⁶⁶ and has been implicated in brain maturation processes (marijuana users showed thinner cortices) in those at risk for schizophrenia.⁷² Notable here is that the potential kindling effect of cannabis on psychotic illness is likely affected by the abovementioned changes in cannabis chemistry. As previously discussed, cannabidiol (CBD)—the component of marijuana that demonstrates antipsychotic properties—is found in smaller concentrations in many recent strains, whereas the percentage of the psychosis-prone component—THC—has increased.^{73,74}

Not only can early, heavy cannabis use potentially contribute to the development and expression of psychotic illness, but ongoing exposure after developing a psychotic disorder can make it worse. In people already suffering with schizophrenia, exposure to THC may lead to an increase in latent psychotic symptoms or relapse back into a psychotic episode.^{64,73} Specifically, patients with psychotic illness who use marijuana, compared to those who do not, tend to have 1) earlier onset of symptoms, 2) more severe and persistent psychotic symptoms, 3) higher relapse rates, and 4) a worse prognosis due to poor treatment adherence.^{70,73} Summarizing the research, Hall et al³⁶ document that cannabis use doubles the risk of developing psychosis from 7 in 1,000 to 14 in 1,000. Therapeutically, however, cessation is salutary: young

people with psychosis who desist from pot have better outcomes, including fewer psychotic symptoms and better social functioning.³⁶

Although the bulk of research on the role of cannabis in severe mental illness is in psychotic disorders, recent research on patients with bipolar illness and posttraumatic stress disorder (PTSD) yields similar findings. Specifically, patients with bipolar disorder who used and then ceased use of marijuana have similar outcomes to those who never used, whereas continued use is associated with greater recurrence risk and functional impairments.⁷⁵ Veterans with PTSD who use marijuana have greater symptom severity of their disorder, use alcohol and other drugs more often, and exhibit more violent behavior than never-users.⁷⁶ Finally, growing evidence of the role of cannabis in other substance use disorders (SUDs) indicates that people with SUDs or who are at risk for developing them are uniquely vulnerable to developing negative effects of cannabis.

Overall, this research suggests that healthcare providers should have targeted risk discussions about cannabis use with adolescents, who are at a higher risk of developing dependence, and individuals at risk for (or suffering from) psychotic illness, bipolar disorder, PTSD, or SUDs.

What about the data on the role of cannabis in other common brain-based disorders (e.g., anxiety, depression), many of which are used by card-carrying medical marijuana users as the reason for their use? Here, unfortunately, research is limited. A few research groups have shown lower perceived quality of life among cannabis users, as well as finding a heightened occurrence of anxiety disorders among cannabis-dependent adults.^{77–79} That said, these studies and others generally indicate that people who only use occasionally to moderately (i.e., who do not qualify for cannabis dependence, or as regular users) generally have the same mental health outcomes as non-users.^{77,78,80}

BAD TO WORSE

Both Aldous Huxley and Jim Morrison famously opined that certain drugs open “the doors of perception.” Does cannabis open the gates of addiction? In short, the “gateway drug” theory posits that the recreational use of “softer” drugs like alcohol, tobacco, and marijuana serves as an easy port of entry into later use of “harder” drugs such as cocaine, heroin, or methamphetamine. The empirical support for this theory largely rests on the observation that most people who develop problems with the latter first experimented with the former. An examination of dairy use, however, exposes this argument’s logical flaw. That is, though many people with opiate dependence have used cannabis prior to developing their heroin habit, a significant majority also ingested milk prior to the onset of their addiction, and yet no one posits a causal connection in this latter case.

Rigorously proving that use of cannabis is consistently associated with a “bad to worse” progression-of-use phenomenon turns out to be methodologically challenging.⁸¹ That said, efforts have been made to answer this question. For example, Olthius et al⁸² looked at the actual circumstances under which people first experimented with a hard drug. This study showed that subjects tended to mix psychoactive substances the first time that they used a new drug. For example, people frequently reported that first-time use of cocaine, heroin, methamphetamine, or a hallucinogen like LSD was in conjunction with marijuana, alcohol, or tobacco, rather than experimenting with the hard drug by itself for the first time.

Additional evidence in support of bad-to-worse causality comes from Agrawal et al,⁴⁶ who performed a twin study examining early cannabis use and later illicit drug use. This study showed a strong association between early cannabis use and later abuse/dependence of other illicit drugs, and—though a large

percentage of the variance in illicit drug use was due to genetic and environmental factors—there was also evidence of a causal influence of early cannabis use. Finally, in a very recent, prospective study (which overcomes methodological limitations inherent in other examinations of this issue), cannabis use during the first sampling period was significantly associated with substance use disorders in a second sampling period three years later.⁸³ In sum, though empirical validation of a direct, causal role of early cannabis use in later addiction to harder drugs is methodologically challenging, and though correlation does not equal causation, several lines of evidence support that the association between early cannabis use and later problems with harder drugs is at least partly due to a causal relationship.^{36,83,84}

How would a gateway process actually work? At a behavioral and interpersonal level, cannabis use likely follows principles of the so-called social contagion seen with alcohol and tobacco,^{85,86} creating “birds of a feather” networks of people with similar behaviors and greater likelihood of exposure to other drug use. At a neurobiological level, animal research points to THC’s ability to change reward circuits in the brain.^{32,33,86} Panlilio et al,⁸⁷ for example, found that exposing rats to THC increased likelihood of self-administration of the most highly addictive drug in humans: nicotine. This effect persisted even when the process to receive the nicotine became more arduous. Interestingly, this effect was not found when a similar experiment was performed with cocaine or heroin in place of nicotine. More recently, Volkow et al³² examined 24 marijuana abusers using a methylphenidate challenge to probe the reactivity of the brain’s dopamine system. They found that compared to normals, marijuana abusers (average 5 joints a day) displayed blunted dopamine responses in key brain areas associated with addiction (i.e., ventral striatum). In a very recent study using an amphetamine

challenge, lower levels of dopamine release in key addiction areas (the striatum) were found in heavy cannabis users; these changes were correlated with inattention, negative symptoms, and poorer working memory.³⁵ These neurobiological differences, researchers hypothesize, may contribute to marijuana abusers’ tendencies to negative emotionality (increased reactivity to stress and irritability) and addictive behaviors.³² Research like this raises the possibility that part of the etiology of marijuana’s bad-to-worse phenomenon is explained by its role in altering brain reward circuits in a way that increases the risk of future addiction.

CONCLUSION

Evaluating the potential harms of a commonly used drug—especially a complex substance like marijuana—is a challenging but vital task. Fully informed awareness of both the potential and proven benefits and the potential and proven harms of marijuana are necessary in order to have rational discussions with patients, teens, and decision makers regarding marijuana use. Based on a review of the current literature, we suggest the mnemonic DDUMB (dependence, driving, underachievement, mental illness, and “bad to worse”) as a tool that captures several of the more well-supported, brain-based risks associated with marijuana. Using this mnemonic, we reviewed five research-supported harms related to marijuana use. First, cannabis dependence (addiction) is real. Second, driving while under the influence of marijuana is unsafe. Third, marijuana use has a strong association with global underachievement. Fourth, marijuana elevates the risk of developing a psychotic illness and worsens the course of several serious mental health conditions in certain individuals. Fifth, though proving causality is complex, evidence supports a “bad to worse” or “gateway” role of cannabis in the

development of other substance use disorders. Important to note, most of these harms are more likely to be present when marijuana use is frequent and starts early (i.e., in adolescence).

Though we don't always heed George Santayana's aphorism about learning from (and therefore being doomed to repeat) our past, a chapter of recent history informs the cannabis conversation. Like a string of white crosses on the shoulder of a dangerous stretch of road, deaths from the recent prescription opiate epidemic stand out as stark examples of the collateral damage from widespread availability of addicting substances and the powerful impact of market forces, medical culture, and societal mores on drug use. Though cannabis is less directly deadly than opiates, all of the factors that buoyed the recent opiate epidemic—availability, economic forces, changing cultural norms—inform the cannabis debate. Healthcare providers, educators, policy leaders, and parents will be well-served by keeping abreast of the burgeoning research on the potential harms of this version of “going green.”

REFERENCES

- Roffman RA. Legalization of marijuana: unraveling quandaries for the addiction professional. *Front Psychiatry*. 2013;4:50.
- Richter KP, Levy S. Big marijuana--lessons from big tobacco. *N Engl J Med*. 2014 Jul 31;371(5):399-401.
- Friese B, Grube JW. Legalization of medical marijuana and marijuana use among youths. *Drugs* 2013;20(1):33-39.
- Schauer GL, King BA, Bunnell RE, et al. Toking, vaping, and eating for health or fun: marijuana use patterns in adults, U.S., 2014. *Am J Prev Med*. 2016;50(1):1-8.
- Atakan Z. Cannabis, a complex plant: different compounds and different effects on individuals. *Ther Adv Psychopharmacol*. 2012;2(6):241-254.
- Monte AA, Zane RD, Heard KJ. The implications of marijuana legalization in Colorado. *JAMA*. 2015;313(3):241-242.
- Paulozzi LJ, Weisler RH, Patkar AA. A national epidemic of unintentional prescription opioid overdose deaths: how physicians can help control it. *J Clin Psychiatry*. 2011;72(5):589-592.
- Walker DD, Stephens R, Roffman R, et al. Randomized, controlled trial of motivational enhancement therapy with nontreatment-seeking adolescent cannabis users: a further test of the teen marijuana check-up. *Psychol Addict Behav*. 2011;25(3):474-484.
- Leung L. Cannabis and its derivatives: review of medical use. *J Am Board Fam Med*. 2011;24(4):452-462.
- Mechoulam R, Parker LA. The endocannabinoid system and the brain. *Annu Rev Psychol*. 2013;64:21-47.
- Gasnier LJ. Reefer Madness. In: Hirliman G (ed). *United States: Motion Picture Ventures*; 1936.
- Van Scoyoc J, Stanger, C, Budney A. Treatment for adolescent marijuana abuse and dependence. *The Brown University Child and Adolescent Behavior Letter*. 2009;25(2):5-7.
- Bachman JG, Johnson LD, O'Malley PM. Explaining recent increases in students' marijuana use: impacts of perceived risks and disapproval, 1976 through 1996. *Am J Public Health*. 1998;88(6):887-892.
- Substance Abuse and Mental Health Services Administration. Results from the 2010 National Survey on Drug Use and Health: Summary of National Findings. January 2012. http://www.ncds.org/images/SAMHSA_Results2010NatlSurveyDrugUseHealth-MentalHealthFindings_1-2012.pdf. Accessed April 1, 2016.
- Morean ME, Kong G, Camenga DR, et al. High school students' use of Eelectronic cigarettes to vaporize cannabis. *Pediatrics*. 2015;136(4):611-616.
- Huddleston T Jr. Legal marijuana sales could hit \$6.7 billion in 2016. *Fortune*. Feb 1, 2016.
- Lovett I. In California, marijuana is smelling more like big business. *New York Times*. April 11, 2016.
- Nelson A. How big is the marijuana market? CNBC2010. April 20, 2010. <http://www.cnbc.com/id/36179677>. Accessed April 2, 2016.
- Chemi E. If the NFL were a real business. *Bloomberg Business*. September 12, 2014. <http://www.bloomberg.com/news/articles/2014-09-12/the-nfl-could-be-a-big-business-with-puny-revenue>. Accessed April 1, 2016.
- Van Zee A. The promotion and marketing of oxycontin: commercial triumph, public health tragedy. *Am J Public Health* 2009;99(2):221-227.
- Kim HS, Monte AA. Colorado cannabis legalization and its effect on emergency care. *Ann Emerg Med*. 2016 Feb 24 [Epub ahead of print]
- ElSohly MA, Mehmedic Z, Foster S, et al. Changes in cannabis potency over the last 2 decades (1995-2014): analysis of current data in the United States. *Biol Psychiatry*. 2016;79(7): 613-619.
- Volkow ND, Baler RD, Compton WM, Weiss SR. Adverse health effects of marijuana use. *N Engl J Med*. 2014;370(23):2219-2227.
- Rehman A. Clearing the haze surrounding state medical marijuana laws: a preemption analysis and proposed solutions. *Houston Journal of Health Law & Policy*. 2014;14.
- Carter GT, Javaher SP, Nguyen MH, et al. Re-branding cannabis: the next generation of chronic pain medicine? *Pain Manag*. 2015;5(1):13-21.
- Grotenhermen F. Pharmacokinetics and pharmacodynamics of cannabinoids. *Clin Pharmacokinet*. 2003;42(4):327-360.
- Allsop DJ, Copeland J, Norberg MM, et al. Quantifying the clinical significance of cannabis withdrawal. *PLoS One*. 2012;7(9):e44864.
- Koob GF, Volkow ND. Neurocircuitry of addiction. *Neuropsychopharmacology*. 2010;35(1):217-238.
- Budney AJ, Moore BA, Vandrey RG,

- Hughes JR. The time course and significance of cannabis withdrawal. *J Abnorm Psychol.* 2003;112(3):393–402.
30. United Nations Office on Drugs and Crime. *World Drug Report 2010.* New York, NY: United Nations; 2010.
31. Hasin DS, Saha TD, Kerridge BT, et al. Prevalence of marijuana use disorders in the United States between 2001–2002 and 2012–2013. *JAMA Psychiatry.* 2015;72(12):1235–1242.
32. Volkow ND, Wang GJ, Telang F, et al. Decreased dopamine brain reactivity in marijuana abusers is associated with negative emotionality and addiction severity. *Proc Natl Acad Sci U S A.* 2014;111(30):E3149–156.
33. Filbey FM, Dunlop J. Differential reward network functional connectivity in cannabis dependent and non-dependent users. *Drug Alcohol Depend.* 2014;140:101–111.
34. Loureiro M, Renard J, Zunder J, Laviolette SR. Hippocampal cannabinoid transmission modulates dopamine neuron activity: impact on rewarding memory formation and social interaction. *Neuropsychopharmacology.* 2015;40(6):1436–1447.
35. an de Giessen E, Weinstein JJ, Cassidy CM, et al. Deficits in striatal dopamine release in cannabis dependence. *Mol Psychiatry.* 2016 Mar 22 [epub ahead of print].
36. Hall W, Degenhardt L. The adverse health effects of chronic cannabis use. *Drug Test Anal.* 2014;6(1–2):39–45.
37. Mason BJ, Crean R, Goodell V, et al. A proof-of-concept randomized controlled study of gabapentin: effects on cannabis use, withdrawal and executive function deficits in cannabis-dependent adults. *Neuropsychopharmacology.* 2012;37(7):1689–1698.
38. Budney AJ, Vandrey RG, Hughes JR, et al. Comparison of cannabis and tobacco withdrawal: severity and contribution to relapse. *J Subst Abuse Treat.* 2008;35(4):362–368.
39. Copersino ML, Boyd SJ, Tashkin DP, et al. Cannabis withdrawal among non-treatment-seeking adult cannabis users. *Am J Addict.* 2006;15(1):8–14.
40. Crean RD, Tapert SF, Minassian A, et al. Effects of chronic, heavy cannabis use on executive functions. *J Addict Med.* 2011;5(1):9–15.
41. Goldstein RZ, Volkow ND. Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry.* 2002;159(10):1642–1652.
42. Chen CY, Storr CL, Anthony JC. Early-onset drug use and risk for drug dependence problems. *Addict Behav.* 2009 Mar;34(3):319–322.
43. Kirisci L, Tarter RE, Ridenour T, et al. Longitudinal modeling of transmissible risk in boys who subsequently develop cannabis use disorder. *Am J Drug Alcohol Abuse.* 2013;39(3):180–185.
44. Shaw P, Eckstrand K, Sharp W, et al. Attention-deficit/hyperactivity disorder is characterized by a delay in cortical maturation. *Proc Natl Acad Sci U S A.* 2007;104(49):19649–19654.
45. Brady JE., Li G. Trends in alcohol and other drugs detected in fatally injured drivers in the United States, 1999–2010. *Am J Epidemiol.* 2014;179(6):692–699.
46. Agrawal A, Neale MC, Prescott CA, Kendler KS. A twin study of early cannabis use and subsequent use and abuse/dependence of other illicit drugs. *Psychol Med.* 2004;34(7):1227–1237.
47. Downey LA, King R, Papafotiou K, et al. The effects of cannabis and alcohol on simulated driving: Influences of dose and experience. *Accid Anal Prev.* 2013;50:879–886.
48. Lenne MG, Dietze PM, Triggs TJ, et al. The effects of cannabis and alcohol on simulated arterial driving: Influences of driving experience and task demand. *Accid Anal Prev.* 2010;42(3):859–866.
49. Hartman RL, Huestis MA. Cannabis effects on driving skills. *Clin Chem.* 2013;59(3):478–492.
50. Buttner A. Review: the neuropathology of drug abuse. *Neuropathol Appl Neurobiol.* 2011;37(2):118–134.
51. Earlywine M. *Understanding Marijuana: A New Look at the Scientific Evidence.* New York: Oxford University Press; 2002.
52. Lachenmeier DW, Rehm J. Comparative risk assessment of alcohol, tobacco, cannabis and other illicit drugs using the margin of exposure approach. *Sci Rep.* 2015;5:8126.
53. Carroll A. Alcohol or marijuana? a pediatrician faces the question. *New York Times.* March 16, 2015. http://www.nytimes.com/2015/03/17/upshot/alcohol-or-marijuana-a-pediatrician-faces-the-question.html?smid=nytcore-iphone-share&smprod=nytcore-iphone&_r=0. Accessed April 1, 2016.
54. Lynskey M, Hall W. The effects of adolescent cannabis use on educational attainment: a review. *Addiction.* 2000;95(11):1621–1630.
55. Bray JW, Zarkin GA, Ringwalt C, Qi J. The relationship between marijuana initiation and dropping out of high school. *Health Econ.* 2000;9(1):9–18.
56. Fergusson DM, Boden JM. Cannabis use and later life outcomes. *Addiction.* 2008;103(6):969–976; discussion 77–78.
57. Meier MH, Caspi A, Ambler A, et al. Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proc Natl Acad Sci U S A.* 2012;109(40):E2657–2664.
58. Daly M. Personality may explain the association between cannabis use and neuropsychological impairment. *Proc Natl Acad Sci U S A.* 2013;110(11):E979.
59. Moffitt TE, Meier MH, Caspi A, Poulton R. Reply to Rogeberg and Daly: no evidence that socioeconomic status or personality differences confound the association between cannabis use and IQ decline. *Proc Natl Acad Sci U S A.* 2013;110(11):E980–E982.
60. Casey BJ, Jones RM, Hare TA. The adolescent brain. *Ann N Y Acad*

- Sci.* 2008;1124:111–126.
61. Jager G, Ramsey NF. Long-term consequences of adolescent cannabis exposure on the development of cognition, brain structure and function: an overview of animal and human research. *Curr Drug Abuse Rev.* 2008;1(2):114–123.
 62. Schweinsburg AD, Nagel BJ, Schweinsburg BC, Park A, et al. Abstinent adolescent marijuana users show altered fMRI response during spatial working memory. *Psychiatry Res.* 2008;163(1):40–51.
 63. Jacobus J, Tapert SF. Effects of cannabis on the adolescent brain. *Curr Pharm Des.* 2014;20(13):2186–2193.
 64. Evins AE, Green AI, Kane JM, Murray RM. The effect of marijuana use on the risk for schizophrenia. *J Clin Psychiatry.* 2012;73(11):1463–1468.
 65. Carey SE, Nestor L, Jones J, et al. Impaired learning from errors in cannabis users: dorsal anterior cingulate cortex and hippocampus hypoactivity. *Drug Alcohol Depend.* 2015 Oct 1;155:175–182.
 66. Zalesky AN, Solowij et al. Effect of long-term cannabis use on axonal fibre connectivity. *Brain.* 2012;135(Pt 7):2245–2255.
 67. Auer R, Vittinghoff E, Yaffe K, et al. Association between lifetime marijuana use and cognitive function in middle age: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *JAMA Intern Med.* 2016;176(3):352–361.
 68. Manrique-Garcia E, Zammit S, Dalman C, et al. Cannabis, schizophrenia and other non-affective psychoses: 35 years of follow-up of a population-based cohort. *Psychol Med.* 2012;42(6):1321–1328.
 69. Gage SH, Hickman M, Zammit S. Association between cannabis and psychosis: epidemiologic evidence. *Biol Psychiatry.* 2016;79(7):549–556.
 70. Patel R, Wilson R, Jackson R, et al. Association of cannabis use with hospital admission and antipsychotic treatment failure in first episode psychosis: an observational study. *BMJ Open.* 2016;6(3):e009888.
 71. Caspi A, Moffitt TE, Cannon M, et al. Moderation of the effect of adolescent-onset cannabis use on adult psychosis by a functional polymorphism in the catechol-O-methyltransferase gene: longitudinal evidence of a gene X environment interaction. *Biol Psychiatry.* 2005;57(10):1117–1127.
 72. French L, Gray C, Leonard G, et al. Early cannabis use, polygenic risk score for schizophrenia and brain maturation in adolescence. *JAMA Psychiatry.* 2015;72(10):1002–1011.
 73. Rey J. Does marijuana contribute to psychotic illness? *Current Psychiatry.* 2007;6(2).
 74. Di Forti M, Marconi A, Carra E, et al. Proportion of patients in south London with first-episode psychosis attributable to use of high potency cannabis: a case-control study. *Lancet Psychiatry.* 2015;2(3):233–238.
 75. Zorrilla I, Aguado J, Haro JM, et al. Cannabis and bipolar disorder: does quitting cannabis use during manic/mixed episode improve clinical/functional outcomes? *Acta Psychiatr Scand.* 2015;131(2):100–110.
 76. Wilkinson ST, Stefanovics E, Rosenheck RA. Marijuana use is associated with worse outcomes in symptom severity and violent behavior in patients with posttraumatic stress disorder. *J Clin Psychiatry.* 2015;76(9):1174–1180.
 77. van Ours JC, Williams J. The effects of cannabis use on physical and mental health. *J Health Econ.* 2012;31(4):564–577.
 78. Lev-Ran S, Le Foll B, McKenzie K, Rehm J. Cannabis use and mental health-related quality of life among individuals with anxiety disorders. *J Anxiety Disord.* 2012;26(8):799–810.
 79. Hasin DS, Kerridge BT, Saha TD, et al. Prevalence and correlates of DSM-5 cannabis use disorder, 2012–2013: findings from the National Epidemiologic Survey on Alcohol and Related Conditions-III. *Am J Psychiatry.* 2016 Mar 4. [Epub ahead of print].
 80. van der Pol P, Liebrechts N, de Graaf R, et al. Mental health differences between frequent cannabis users with and without dependence and the general population. *Addiction.* 2013;108(8):1459–1469.
 81. Kandel DB. Does marijuana use cause the use of other drugs? *JAMA.* 2003;289(4):482–483.
 82. Olthuis JV, Darredeau C, Barrett SP. Substance use initiation: the role of simultaneous polysubstance use. *Drug Alcohol Rev.* 2013;32(1):67–71.
 83. Blanco C, Hasin DS, Wall MM, et al. Cannabis use and risk of psychiatric disorders: prospective evidence from a US national longitudinal study. *JAMA Psychiatry.* 2016 Feb 17 [epub ahead of print].
 84. Moss HB, Chen CM, Yi HY. Early adolescent patterns of alcohol, cigarettes, and marijuana polysubstance use and young adult substance use outcomes in a nationally representative sample. *Drug Alcohol Depend.* 2014;136:51–62.
 85. Christakis NA, Fowler JH. The collective dynamics of smoking in a large social network. *N Engl J Med.* 2008 May 22;358(21):2249–2258.
 86. Rosenquist JN, Murabito J, Fowler JH, Christakis NA. The spread of alcohol consumption behavior in a large social network. *Ann Intern Med.* 2010;152(7):426–433, W141.
 87. Panlilio LV, Zanettini C, Barnes C, et al. Prior exposure to THC increases the addictive effects of nicotine in rats. *Neuropsychopharmacology.* 2013;38(7):1198–1208. ■